

INFRARED SPECTROSCOPY OF HYDROGEN CYANIDE IN SOLID PARAHYDROGEN (BRIEFING CHARTS)

C. Michael Lindsay,
National Research Council, Post Doctoral Research Associate
Mario E. Fajardo
Air Force Research Laboratory
Munitions Directorate
AFRL/MNME
Eglin AFB, FL 32542-6810



JULY 2006

CONFERENCE PAPER

This paper will be presented at the 61st Ohio State University International Symposium on Molecular Spectroscopy, Ohio State University, Columbus, OH, 19-23 June 2006. One of the authors is a U.S. Government employee working within the scope of his position; therefore, the U.S. Government is joint owner of the work. This paper will be published in the proceedings and the publisher may assert copyright. If so, the Government has the right to copy, distribute, and use the work. Any other form of use is subject to copyright restrictions.

This paper is published in the interest of the scientific and technical information exchange. Publication of this paper does not constitute approval or disapproval of the ideas or findings.

DISTRIBUTION A: Approved for public release; distribution unlimited.
Approval Confirmation #AAC/PA 06-02-0-271.

AIR FORCE RESEARCH LABORATORY, MUNITIONS DIRECTORATE

■ Air Force Materiel Command ■ United States Air Force ■ Eglin Air Force Base

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE Conference Paper		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Infrared Spectroscopy of Hydrogen Cyanide In Solid Parahydrogen (BRIEFING CHARTS)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Mario E. Fajardo C. Michael Lindsay (National Research Council, Post Doctoral Research Associate) assigned to the Air Force Research Laboratory, Munitions Directorate, AFRL/MNME				5d. PROJECT NUMBER 2502	
				5e. TASK NUMBER 10	
				5f. WORK UNIT NUMBER 15	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory, Munitions Directorate AFRL/MNME Eglin AFB, FL 32542-6810				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-MN-EG-TP-2006-7403	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Same as Block 7				10. SPONSOR/MONITOR'S ACRONYM(S) AFRL-MN-EG	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) Same as Block 8	
12. DISTRIBUTION / AVAILABILITY STATEMENT DISTRIBUTION A: Approved for public release; distribution unlimited. Approval Confirmation #AAC/PA 06-02-06-271					
13. SUPPLEMENTARY NOTES This paper will be presented at the 61st Ohio State University International Symposium on Molecular Spectroscopy, Ohio State University, Columbus, OH, 19-23 June 2006. One of the authors is a U.S. Government employee working within the scope of his position; therefore, the U.S. Government is joint owner of the work. This paper will be published in the proceedings and the publisher may assert copyright. If so, the Government has the right to copy, distribute, and use the work. Any other form of use is subject to copyright restrictions.					
14. ABSTRACT <u>Solid parahydrogen features:</u> <ul style="list-style-type: none"> Weak intermolecular interactions Slow relaxation timescales Quantum crystal: self-annealing via tunneling Homogeneous environment Large intermolecular distance, 3.78 Å <u>Parahydrogen as a matrix:</u> <ul style="list-style-type: none"> Very small matrix shifts, predictable (-0.5 %) Nearly free rotation for small molecules High impurity mobility Narrow spectral linewidths (<100 MHz possible) Dopants induce infrared activity in matrix 					
15. SUBJECT TERMS Solid Parahydrogen, Hydrogen Cyanide, Infrared Spectroscopy					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 20	19a. NAME OF RESPONSIBLE PERSON Mario E. Fajardo
a. REPORT UNCLASSIFIED	b. ABSTRACT UNCLASSIFIED	c. THIS PAGE UNCLASSIFIED			19b. TELEPHONE NUMBER (include area code)

Infrared Spectroscopy Of Hydrogen Cyanide In Solid Parahydrogen



**Dr. C. Michael Lindsay
& Dr. Mario E. Fajardo
AFRL/MNME**

DISTRIBUTION A: Approved for public release; distribution unlimited. AAC/PA 06-02-06-271

61th International Symposium on Molecular Spectroscopy, Talk RD10, 22 June 2006, The Ohio State University, Columbus, OH

This paper is published in the interest of the scientific and technical information exchange. Publication of this paper does not constitute approval or disapproval of the ideas or findings.



Infrared Spectroscopy Of Hydrogen Cyanide In Solid Parahydrogen



C. Michael Lindsay and Mario E. Fajardo

AFRL/MNME, Energetic Materials Branch, Ordnance Division,
U.S. Air Force Research Laboratory, 2306 Perimeter Rd, Eglin
AFB, FL 32542-5910 c.michael.lindsay@eglin.af.mil




Solid parahydrogen



Solid parahydrogen features:

- Weak intermolecular interactions
- Slow relaxation timescales
- Quantum crystal: self-annealing via tunneling
- Homogeneous environment
- Large intermolecular distance, 3.78 Å




Parahydrogen as a matrix:

- Very small matrix shifts, predictable (-0.5 %)
- Nearly free rotation for small molecules
- High impurity mobility
- Narrow spectral linewidths (<100 MHz possible)
- Dopants induce infrared activity in matrix




Rotation of Molecules in Solid Parahydrogen



Data from the groups of M. E. Fajardo, Y.-P. Lee, and T. Momose (1995-2006).




Why study HCN?



* *D. T. Moore, M. Ishiguro, and R. E. Miller, JCP 115, 5144 (2001)*



H₂-HCN Potential



If solid parahydrogen behaved classically, the HCN will not rotate...


CCSD(T)/aug-cc-pVTZ calculations D. T. Moore, M. Ishiguro, and R. E. Miller, JCP 115, 5144 (2001)

61th International Symposium on Molecular Spectroscopy, Talk RD10, 22 June 2006, The Ohio State University, Columbus, OH

Approved for Public Release; Distribution Unlimited. AAC/PA 06-02-06-271




Rapid vapor deposition of $p\text{-H}_2$



- Reflection based set-up (FTIR)
- $[o\text{-H}_2] < 100$ ppm
- $p\text{-H}_2$ deposition rate ~ 150 mmol/hr $\rightarrow \sim 1$ mm/hr thickness
- Deposition at 2.3 K. Sample annealed at 4.3 K for 30 min.



Spectrum of annealed solid $p\text{-H}_2$ with and without HCN




61th International Symposium on Molecular Spectroscopy, Talk RD10, 22 June 2006, The Ohio State University, Columbus, OH

Approved for Public Release; Distribution Unlimited. AAC/PA 06-02-06-271




Spectrum of annealed solid p -H₂ with and without HCN





HCN ν_1 region







Assignment tools




Temperature Dependence



Orthohydrogen Dependence



Combination Differences





Spectrum was interpreted by its:

- Temperature dependence
- Orthohydrogen concentration dependence
- Transition frequency combination differences
- **Polarization dependence!**



Polarization spectroscopy





Polarization analysis provides:

- An assignment of the perpendicular and parallel components to the crystal field perturbed rotational structure (*i.e.* Δm)
- $Abs_{s-pol} - Abs_{p-pol} = \frac{1}{2} \mu^2_{\perp} - \frac{1}{2} \mu^2_{\parallel}$
- A measures of the fraction of HCP lattice with c -axis normal to substrate:




An aside... new insights into annealing mechanism in $p\text{-H}_2$



- Before annealing, less than 5% of the HCP features are oriented normal to the substrate.
 - After annealing, ~80% of the HCP features are oriented normal to substrate!!
- Upon deposition, HCP lattices randomly oriented, but upon annealing gain a macroscopic orientation to the laboratory frame.



Interpretation of the fine structure





Interpretation of the fine structure in the HCN ν_1 fundamental region




- All peaks assigned!
- Each transition fit to a Lorentzian profile. Standard deviation of the fit for the last digits is shown in parentheses.
- Polarization dependence was not as clear for the higher order clusters.

Species	$ J',M'\rangle \leftarrow J'',M''\rangle$	Polarization	ν_0
HCN	$ 1,1\rangle \leftarrow 2,2\rangle$	\perp	3297.989(5)
HCN	$ 1,1\rangle \leftarrow 2,1\rangle$	\parallel	3298.678(3)
HCN	$ 1,0\rangle \leftarrow 2,1\rangle$	\perp	3299.285(10)
HCN	$ 0,0\rangle \leftarrow 1,0\rangle$	\parallel	3300.282(1)
HCN	$ 0,0\rangle \leftarrow 1,1\rangle$	\perp	3300.764(1)
HCN	$ 1,1\rangle \leftarrow 1,1\rangle$	\perp	3302.880(1)
HCN	$ 1,1\rangle \leftarrow 0,0\rangle$	\perp	3305.105(1)
HCN	satellite	\perp	3305.530(2)
HCN	$ 1,0\rangle \leftarrow 0,0\rangle$	\parallel	3305.709(1)
HCN	$ 2,1\rangle \leftarrow 1,0\rangle$	\perp	3306.572(1)
HCN	$ 2,1\rangle \leftarrow 1,1\rangle$	\parallel	3307.055(1)
HCN	$ 2,2\rangle \leftarrow 1,1\rangle$	\perp	3307.778(1)
HCN	R(2)?	\parallel	3309.328(16)
HCN	$ 2,2\rangle \leftarrow 0,0\rangle$	\perp	3310.001(5)
HCN-oH2 (ip)	in-plane cluster	\perp	3302.708(1)
HCN-oH2 (op)	out-of-plane cluster	\parallel	3302.946(1)
oH2-HCN (ip)	in-plane cluster	\perp	3308.046(1)
oH2-HCN (op)	out-of-plane cluster	\parallel	3309.803(1)
(oH2)n-HCN			3302.238(1)
(oH2)n-HCN			3302.405(1)
(oH2)n-HCN			3302.474(1)
(oH2)n-HCN			3302.579(1)
(oH2)n-HCN			3302.663(1)
(oH2)n-HCN			3302.760(1)
(oH2)n-HCN			3310.775(7)
(oH2)n-HCN			3310.859(25)
(oH2)n-HCN			3311.128(5)
(oH2)n-HCN			3311.479(12)
(oH2)n-HCN			3311.717(9)
(HCN)n			3297.771(11)
(HCN)n			3297.844(4)
(HCN)n			3298.697(4)
(HCN)n			3298.739(4)



Interpretation of the fine structure



Crystal field theory (linear molecule in HCP lattice) *

$$H = H_{r,v} + V_{\text{cry}}$$

$$V_{\text{cry}} = \epsilon_2 C_{2,0}(\Omega_{\text{HCN}}) + \epsilon_3 [C_{3,-3}(\Omega_{\text{HCN}}) - C_{3,3}(\Omega_{\text{HCN}})]$$

$$\text{where } C_{l,m}(\Omega) = \left(\frac{4\pi}{2l+1} \right)^{1/2} Y_{l,m}(\Omega)$$

“Fitting” Results: (10 levels, 9 parameters)

	<i>p</i> -H ₂	Gas
<i>B</i>	1.475(5)	1.478
ΔB	-0.005(5)	-0.010
<i>D</i>	0.0675(5)	2.910×10^{-6}
ΔD	0.0032(5)	0.025×10^{-6}
ν_0	3302.989(5)	3311.4770
ϵ_2	-1.274(5)	-
$\Delta \epsilon_2$	-0.193(5)	-
ϵ_3	6.850(5)	-
$\Delta \epsilon_3$	0.767(5)	-

Identical!

$\times 20,000!$

very large!

NOTE: Above perturbation does not converge for large *D*!
Treatment was modified to incorporate centrifugal distortion after crystal field calculation to avoid this problem.

* Perturbation terms determined by T. Momose, unpublished.



Summary



- **AFRL matrix isolation spectroscopy lab back in business (after 5 years!)**
- **Polarization spectroscopy is helpful with reflection configuration**
 - 1) New insights into annealing process
 - 2) Powerful assignment tool
- **Despite size, and strong interaction, HCN does rotate in HCP p -H₂**
 - 1) Large crystal field perturbation (perturbation theory may not be appropriate)
 - 2) Large effective centrifugal distortion constant, (D), but B unchanged!
 - 3) Crystal field calculation does not converge properly when D is large
 - 4) Incorporating D after the crystal field calculation works well
 - 5) A single o -H₂ inhibits rotation of HCN
- **Other interesting features left out of the talk...**
 - 1) Rotational excitation lifetime is sensitive to HCN concentration
 - 2) Strong induced IR activity in H₂ fundamental region
 - 3) HCN cluster formation



Molecular rotation in solid p -H₂...

What should be next

